


Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

<p>CAS # 375-85-9</p> 	<p>Perfluoroheptanoic acid Synonym¹s: Tridecafluoro-1-heptanoic acid; Tridecafluoroheptanoic acid; HSDB 8293 RTECS #²: MJ2415000 EINECS #³: 206-798-9 Molecular Weight⁴: 364.0569 Molecular Formula⁵: C7-H-F13-O2</p> <p>Related CAS #'s: 20109-59-5 (hydrochloride salt); 6130-43-4 (ammonium salt)</p>
PHYSICAL CHARACTERISTICS	
<i>Primary Use</i>	Long-chain perfluoroalkancarboxylic acids and their salts are surface-active chemicals (surfactants), which greatly reduce the surface tension (surface energy) of water, aqueous solutions, and organic liquids even at low concentrations. These acids (C6- C12) and derivatives are used as wetting, dispersing, emulsifying, and foaming agents. /Long-chain perfluoroalkancarboxylic acids/ ⁶
<i>Physical state, odor at room temperature & pressure</i>	Beige crystalline solid ⁷ ; Low melting solid ⁸
<i>Melting point; Boiling point</i>	30 deg C ⁹ ; 175 deg C ¹⁰
<i>Solubility</i>	In water, 3.65 mg/L at 25 deg C (est) ¹¹
<i>Specific Gravity</i>	1.792 g/cu cm at 20 deg C/4 deg C ¹²
SAFETY/PHYSICAL HAZARDS	
<i>Vapor Pressure</i>	0.133 mm Hg at 25 deg C ¹³
<i>Flammability</i>	Not found
<i>Flashpoint</i>	> 113.00 deg C (> 235.40 deg F) - closed cup ¹⁴
<i>Flammability Rating</i>	Not available
<i>Auto Ignition Point</i>	Not found
<i>Combustion products</i>	Carbon oxides, Hydrogen fluoride ¹⁵
<i>Explosivity (UEL, LEL, shock sensitive)</i>	Not found
<i>Oxidizer</i>	Not found
<i>Corrosivity</i>	
<i>pH</i>	pKa = -2.29 (est) ¹⁶
<i>Reactivity</i>	Incompatible materials: Strong oxidizing agents ¹⁷
<i>Viscosity</i>	Not found
<i>Odor Threshold</i>	Not found
<i>Particle size, shape, respirable fraction</i>	Not found
<i>Other physical hazards associated</i>	Not found

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

<i>with process: Heat, gases under pressure, noise, vibration, ergonomic hazard</i>	
HEALTH HAZARDS	
Acute Toxicity	
<i>Oral LD₅₀</i>	"An acute lethal dose of 670 mg/kg bw has been reported for PFHpA. Rats were administered the undiluted liquid in doses ranging from 300 to 3,375 mg/kg. Necrosis of the stomach was observed in animals which died (unpublished study provided by Dupont Co.)." ¹⁸
<i>Dermal LD₅₀</i>	Not found
<i>Inhalation LC₅₀</i>	Not found
<i>Intraperitoneal LD₅₀</i>	Not found
Chronic or Sub-chronic Toxicity	
<i>IARC rating</i>	Not found in the IARC database as of 05/17/18
<i>Carcinogenicity</i>	Not found in the CCRIS database
<i>Neurotoxicity</i>	Not found in Haz-Map
<i>Developmental/Reproductive Toxicity</i>	<p>Not found on the Prop 65 List as of 05/17/18</p> <p>"Plasma concentrations of perfluoroheptanoic acid (PFHpA), perfluorohexane sulfonic acid (PFHxS) and perfluorononanoic acid (PFNA) were inversely associated with endometriosis-related infertility, but the associations were attenuated in the sensitivity analyses (Wang et al 2017)."¹⁹</p> <p>See Ecological Toxicity box below – Kim et al 2015</p>
<i>Genotoxicity/Mutagenicity</i>	<p>"This is the first report of the effects of perfluorinated acids on the activity of purified enzymes. The results show these compounds have a very low acute biological activity. The observed effective concentrations lie in the millimole range, which is well above probable intracellular concentrations. A relationship was found between the toxicity of the perfluorinated carboxylic acids and the perfluorocarbon chain length: in every test system applied, the longer the perfluorocarbon chain, the more toxic was the acid. The lowest effective concentrations were thus recorded for perfluorononanoic and perfluorodecanoic acids (Mulkiewicz et al 2007)."²⁰</p> <p>"A chain length-EC₅₀ dependence has been clearly observed. Estimated values of EC₅₀ decreased with elongation of fluorocarbon chain (PFHxA > PFHpA > PFOA > PFNA > PFDA > PFDoA > PFTeDA). Further elongation (C16 and C18) did not deepen the effect but even partially reversed it. The effect was intensified after longer exposure (72 h);... (Kleszczyński 2007)."²¹</p>

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

	<p>“The results of our study clearly demonstrate that inductions of stress responsive genes occur for the different compounds and confirm some of the known mechanisms of work for well studied compounds like PFOA and PFOS, and in addition provide new information for less studied compounds. Few inductions were observed after exposure to the low carbon number carboxylic acids, ... PFHpA (CF₃(CF₂)(₅)C(O)O(-)) at equimolar concentrations (0.0156-1 mM). Overall, the effects seen at the level of gene expression were higher for the sulfonic acids than for the carboxylic acids, but the effect of the chain length is more important than the effect of the functional group (Nobels et al 2010).”²²</p>
<i>Endocrine Disruption</i>	Added to TEDX List 5/5/11 (information updated 12/6/17)
<i>Thyroid</i>	Bloom 2010 – No values exceeded the LOD, so the substance was not considered further in this paper.
<i>Immunotoxicity</i>	Nothing found as of 05/17/18
<i>Other organ toxicity</i>	<p>“Wolf et al. (2012) showed an increase in activity of human and mouse PPARα, in vitro with PFHpA (NICNAS 2018)”²³</p> <p>“Various studies have investigated the effects of PFHpA on liver enzymes. The maximum induction of peroxisomal beta-oxidation was assessed in Wistar rats (Kudo et al., 2000). A slight, but significant increase in liver beta-oxidation was induced by treatment with 160 mg/kg PFHpA, while a significant increase in liver beta-oxidation was detected following 10 mg/kg PFOA. The potency of induction of hepatomegaly, peroxisomal beta-oxidation and microsomal 1-acylglycerophosphocholine acyltransferase in the liver was investigated in mice (Kudo et al., 2006). Similarly to all PFCA tested, PFHpA induced hepatomegaly and peroxisomal beta-oxidation. The potency of these effects is reported to increase with increasing chain length. The doses required to induce an approximately 1.7-fold increase in liver weight relative to body weight in male mice were >150, 50, 5, and 5 mg/kg for PFHxA, PFHpA, PFOA and perfluorononanoic acid (PFNA), respectively. Fischer-344 (F344) rats treated with PFHpA (150 mg/kg) showed no increase in hepatic peroxisomal fatty acyl CoA-oxidase (FAO) activity (an indicator of peroxisome induction) on days three or five post-treatment (Goecke-Flora & Reo, 1996). In Wistar rats, the activity of peroxisomal beta-oxidation was not affected at doses up to 30 mg/kg body weight in either male or female rats (Kudo & Kawashima, 2003). Peroxisomal beta-oxidation induction is considered to depend only on the number of PFCA molecules in hepatocytes, rather than the difference in their chemical structures (Kudo & Kawashima, 2003; Kudo et al., 2006)(NICNAS 2018)”²⁴</p>

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

Skin, Eye and Respiratory Effects	
<i>Irritant – Skin, Eye, or Respiratory</i>	“...skin irritant and/or corrosive effects cannot be ruled out for PFHpA”(NICNAS 2018). ²⁵
<i>Corrosive – S, E, or R</i>	Corrosive as concentrated acid; pKa = -2.29 (est) ²⁶
<i>Permanent Damage – S, E, or R</i>	Not found
<i>Sensitizer– S & R</i>	Not found in AOEC database
<i>Asthmagen – Initiator or Exacerbator</i>	Not found in AOEC database
<i>Skin Absorption, Kp</i>	Not found
<i>LOAEL</i>	
<i>NOAEL</i>	
<i>Benchmark Dose Response (BMD)</i>	
<i>Toxicokinetics</i>	<p>“5:3 acid [F(CF₂)₅CH₂CH₂COOH] was the most abundant metabolite for 6:2 FTI (3.3-6.3 mol%) and 6:2 FTOH (9-12 mol%). Perfluorobutanoic acid (PFBA), perfluoropentanoic acid (PFPeA), and perfluorohexanoic acid (PFHxA) in sum accounted for 1.3-2.2 mol% from 6:2 FTI and 2.7-4.4 mol% from 6:2 FTOH biotransformation. Perfluoroheptanoic acid (PFHpA) accounted for 0.14-0.36 mol% from 6:2 FTI but only 0.01-0.06 mol% from 6:2 FTOH biotransformation. These results suggest that mammalian systems exposed to 6:2 FTI or 6:2 FTOH would form 5:3 acid, PFBA, PFPeA, PFHxA as the primary stable metabolites, whereas more PFHpA would be expected from 6:2 FTI biotransformation (Ruan et al 2014).”²⁷</p> <p>“In a one-day inhalation study of 6:2 FTOH in rats, PFBA, PFHxA, PFHpA and 5:3 Acid were determined to be the major metabolites in plasma with calculated elimination half-lives of 1.3-15.4h and metabolic yields up to 2.7 mol%. In five-day and 23-day inhalation studies and a 90-day oral study of 6:2 FTOH, the plasma or serum concentration profile of 5:3 Acid was several-fold higher than concentrations observed in the single day study, resulting in an estimated elimination half-life of 20-30 d. In contrast, the concentrations of PFBA, PFHxA and PFHpA showed little or no concentration increase with repeated exposure. Elimination half-lives of PFHxA, PFHpA and 5:3 Acid in humans were estimated from a study of professional ski wax technicians who were occupationally exposed to aerosolized and volatilized components of fluorinated glide wax. The resulting human elimination half-life values of PFHxA, PFHpA and 5:3 Acid were 32, 70 and 43 d, respectively. Based on a one compartment toxicokinetic model, current environmental air concentrations of 6:2 FTOH are estimated to result in plasma concentrations of PFHxA, PFHpA and 5:3 Acid that are less than or equal to typical LOQ values, in agreement with extant biomonitoring</p>

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

	<p>results (Russell et al 2015).”²⁸</p> <p>“The mouse experiment showed that short-chained PFCAs (C6 and C7) were rapidly eliminated in the urine, whereas long-chain PFCAs (C8 to C14) accumulated in the liver and were excreted slowly in feces. Urinary clearance of PFCAs in humans also decreased with increasing alkyl chain lengths, while biliary clearances increased. C9 to C10 had the smallest total clearance for both mice and humans. However, disparities existed in the magnitude of the total clearance between mice and humans. A slightly higher partition ratio (brain/serum) was observed for long-chained PFCAs in mice, but this was not observed in the corresponding partition ratio in humans (CSF/serum) (Fujii et al 2015).”²⁹</p> <p>“The half-life for PFHpA in female and male rats was reported as 1.2 and 2.4 hours, respectively, compared with 5.63 and 0.08 days for PFOA (Ohmori et al., 2003).”³⁰</p> <p>“In studies in mice and rats, PFHpA was shown to be rapidly eliminated in the urine. Total clearance in animals and humans decreases with increasing chain length, with the clearance rate for PFHpA higher than PFOA, but lower than PFHxA (Han et al., 2012; Fujii et al. 2015). Following intraperitoneal (i.p.) injection in Wistar rats, 92 % of a PFHpA dose was rapidly eliminated in urine within 120 hours after dosing (Kudo et al., 2001). Similar results were seen in NJcl mice administered PFHpA intravenously (i.v.) or by gavage. When administered intravenously, the chemical was rapidly and almost completely eliminated in the urine after 24 hours (99 % for males and 66 % for females) and only a small amount was excreted in faeces (3 % for males and 13 % for females). Gavage-administered animals showed similar clearance patterns to i.v. administered animals (Fujii et al., 2015)(NICNAS 2018)”.³¹</p>
<i>Metabolites</i>	
<i>Synergistic or Antagonistic Effects</i>	
Environmental and Human Health Exposure and Risk Values	
<i>RfC/RfD</i>	Not in IRIS database as of 05/18/18
<i>ATSDR-MRL</i>	Not on list as of 05/18/18
<i>Adverse Effect Levels: DNEL, PNEC, PNEL</i>	
Health Based Exposure Limits	
<i>NIOSH-REL/IDLH/Ceiling Limits</i>	None exist
<i>OSHA-PEL</i>	None exist

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

<i>ACGIH TLV-TWA</i>	None exist
<i>TLV-STEL</i>	None exist
<i>Biomonitoring Action Limits</i>	Connecticut DPH set a drinking water Action Level for private wells in 2016 for PFAS that is the same as the EPA Health Advisory (70 ppt) but has added three additional PFAS (PFNA, PFHxS, PFHpA) to the group. The sum of this group of 5 PFAS must be below the target concentration of 70 ppt. These additional PFAS have produced some of the same health effects as PFOS and PFOA (CT DPH 2017). ³²
<i>Drinking Water Standards</i>	
<i>Other</i>	Monitoring requirements for unregulated contaminants: Contaminant, perfluoroheptanoic acid; Minimum reporting level, 0.01 ug/L; Period during which monitoring to be completed, 1/1/2013-12/31/2015. ³³ ECHA CLP Database (non-harmonized classifications): H302 – Acute Tox. 4; H314 – Skin Corr. 1B; H290 - Met. Corr. 1; H314 – Skin Corr. 1C; H318 – Eye Dam. 1. ³⁴
ENVIRONMENTAL & ECO-SYSTEM HAZARDS	
PBT	<p>Presence in the environment:</p> <p>“PFOS, PFOA, PFHxS, PFHpA [Median concentration in dust: 10 – 97.3 ng/g] and PFNA were found in all of the vacuum dust samples and dust from eight homes contained all 16 PFCs included in our analysis (Knobeloch et al 2012).”³⁵</p> <p>“Among samples with detectable PFAS levels, each additional military site within a watershed's eight-digit hydrologic unit is associated with a 20% increase in PFHxS, a 10% increase in both PFHpA and PFOA, and a 35% increase in PFOS (Hu et al 2016).”³⁶</p>
<i>Bioaccumulation</i>	<p>Perfluorinated chemicals have been observed to bioaccumulate by binding to proteins in plasma and liver, rather than the more conventional partitioning to fatty tissue (Ng and Hungerbuhler 2014).³⁷</p> <p>“Ericson et al. (2008b) analyzed 14 PFCs in drinking water (tap and bottled) from Tarragona Province (Catalonia, Spain). In 2007, municipal drinking (tap) water was obtained in public fountains of the three most populated towns in Tarragona Province. The bottled water samples from four commercial companies, whose water spring has different origins, were purchased from a supermarket. This is, to the best of our knowledge, the first study in which bottled water was analyzed. The PFC levels in tap water varied among the four places. In the Valls sample, the highest PFC levels were found; PFHpA (3.02 ng/L), PFOS</p>

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

(0.44 ng/L), and PFOA (6.28 ng/L)(D'Hollander et al 2010).”³⁸

“Perfluoroalkyl and polyfluoroalkyl substances (PFASs) are commonly used in industrial applications and consumer products, and their potential health impacts are of concern, especially for vulnerable population like fetuses. However, in utero exposure to PFASs and health implications are far from fully characterized in China. To fill in the gap, we analyzed 10 PFASs in cord plasma samples (N=687) collected in Shanghai between 2011 and 2012, one of the regions widely polluted with PFASs in China. A questionnaire survey on maternal and diet-related factors was conducted. Except for perfluoroheptanoic acid (PFHpA) [detected in 30% of the samples] and perfluorooctane sulfonamide (PFOSA), all other PFASs were detected in >90% of the samples (Wang et al 2016c).”³⁹

Percent detection and level of PFHpA in:

U.S. NHANES (ages 12-19)(various years from 1999-2010 – see Table 6-16 in ATSDR draft profile): 6.2 (Geometric mean <0.3 ng/mL) - 10%.

**Percent detection not specified for the adolescent age group samples. Percentage listed here is for the total sample population.

U.S. children (ages 2-12)(1994-1995): Not detected above LLOQ.

Umbilical Cord Serum (Baltimore): 2% >LOD, Min = <0.4 ng/mL, Max= 2.6 ng/mL. (Japan): Not detected.

Breast Milk (MA): <1% >LOQ, Min = <0.010 ng/mL, Max = 0.0234 ng/mL. Not detected in study from Sweden.

Maternal Serum (China): 37% > LOD, Min = <0.005 ng/mL, Max = 0.0067 ng/mL. Not detected in study from Sweden.

For additional information see ATSDR draft profile pages 411-413 (ATSDR 2015).⁴⁰

“PFHpA was detected in pizza and microwave popcorn at 1.5–2.0 ng/g, wet weight. (ATSDR 2015)”⁴¹

“there is evidence that PFHpA bioaccumulates in earthworms (Zhao et al., 2013[c]) and humans (Freberg et al., 2010, Nilsson et al., 2010) with a geometric mean elimination half-life in ski-wax technicians of 98 day (range 23 to 599 days). In addition, it has also been shown that PFHpA is maternally transferred in polar bears (Bytingsvik et al., 2012) and

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

	<p>humans (Zhang et al., 2013b). In contrast, trophic magnification of PFHpA was investigated but not observed in an arctic (Kelly et al., 2009), a subtropical pelagic (Loi et al., 2011), nor a remote terrestrial (Muller et al., 2011) food web. Also, there are human maternal transfer studies that did not detect PFHpA in maternal and/or cord blood (Monroy et al., 2008, Kim et al., 2011, Gutzkow et al., 2012). Thus, data from public literature on the bioaccumulation potential of PFHpA are equivocal, and at this moment it is too premature to conclude upon the B-criterion.” (ECHA 2015a)⁴²</p> <p>In earthworm study, $BSAF_{kinetic} = 0.122 \pm 0.006$ (g_{oc}/g_{dw}); BSAF measured based on wet wt of organisms at three exposure concentrations: at 100 ng/g: 0.040 ± 0.023; at 200 ng/g: 0.019 ± 0.002; at 500 ng/g: 0.008 ± 0.001. "According to our study, PFHxA, PFHpA, PFOA, PFBS and PFHxS, which have seven or less fluorinated carbons, displayed distinct bioaccumulative ability. This could be due to the active ingestion of soil through the gut and the high protein content of the earthworms. (Zhao et al 2013)⁴³</p> <p>In Zhang, et al, 2013, a coefficient, T_{MB-CB} was calculated to indicate maternal transfer efficiency of each PFAS between maternal blood and cord blood. A U-shaped trend was observed for PFCAs: “the highest T_{MB-CB} among all the measured PFCAs was for PFHpA (C7). A descending trend in T_{MB-CB} was observed with the increasing carbon chain length from PFHpA (C7) to PFDA (C10); the T_{MB-CB} reached a minimum value for PFDA (C10, $T_{MB-CB} = 0.25$), and then increased with the increasing carbon chain length from PFDA (C10) to PFDoDA (C12)”.⁴⁴</p> <p>Phytoaccumulation – PFHpA, PFOA, PFNA and PFBS were taken up by yam root, maize cob and sugarcane stem (Dalahmeh et al 2018).⁴⁵</p> <p>“In blood of the Gdańsk Coast inhabitants...PFHpA, were found (Falandysz 2006 – Article in Polish).”</p>
BAF	
BCF	
BMF	
Ecological Toxicity	<p>“The present study assessed the transactivation potencies of the Baikal seal (<i>Pusa sibirica</i>) peroxisome proliferator-activated receptor α (BS PPARα) by perfluorochemicals (PFCs) having various carbon chain lengths (C4-C12) using an in vitro reporter gene assay. Among the twelve PFCs treated with a range of 7.8-250 μM concentration, eight perfluoroalkyl carboxylates (PFCAs) and two perfluoroalkyl sulfonates</p>

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

(PFSAs) induced BS PPAR α -mediated transcriptional activities in a dose-dependent manner. To compare the BS PPAR α transactivation potencies of PFCs, the present study estimated the PFOA induction equivalency factors (IEFs), a ratio of the 50% effective concentration of PFOA to the concentration of each compound that can induce the response corresponding to 50% of the maximal response of PFOA. The order of IEFs for the PFCs was as follows: PFOA (IEF: 1)>PFHpA (0.89)>PFNA (0.61)>PFPeA (0.50)>PFHxS (0.41)>PFHxA (0.38) \approx PFDA (0.37)>PFBA (0.26)=PFOS (0.26)>PFUnDA (0.15)>>PFDoDA and PFBuS (not activated). The structure-activity relationship analysis showed that PFCAs having more than seven perfluorinated carbons had a negative correlation ($r=-1.0$, $p=0.017$) between the number of perfluorinated carbons and the IEF of PFCAs, indicating that the number of perfluorinated carbon of PFCAs is one of the factors determining the transactivation potencies of the BS PPAR α . The analysis also indicated that PFCAs were more potent than PFSAs with the same number of perfluorinated carbons. Treatment with a mixture of ten PFCs showed an additive action on the BS PPAR α activation. Using IEFs of individual PFCs and hepatic concentrations of PFCs in the liver of wild Baikal seals, the PFOA induction equivalents (IEQs, 5.3-58 ng IEQ/g wet weight) were calculated. The correlation analysis revealed that the hepatic total IEQs showed a significant positive correlation with the hepatic expression levels of cytochrome P450 4A-like protein ($r=0.53$, $p=0.036$). This suggests that our approach may be useful for assessing the potential PPAR α -mediated biological effects of complex mixtures of PFCs in wild Baikal seal population (Ishibashi et al 2011).⁴⁶

“Whole-mount in situ hybridization, reverse transcriptase-polymerase chain reaction (RT-PCR), and histologic analyses detected severe defects in the liver and heart following exposure to PFHxA or PFHpA. In addition, immunoblotting revealed that PFHpA significantly increased the phosphorylation of extracellular signal-regulated kinase (ERK) and c-Jun N-terminal kinase (JNK), while PFHxA slightly increased these, as compared with the control. These results suggest that PFHxA and PFHpA are developmental toxicants and teratogens, with PFHpA producing more severe effects on liver and heart development through the induction of ERK and JNK phosphorylation (Kim et al 2015).⁴⁷

“Earthworms were exposed to spiked soil for 21 days. Concentrations of these compounds in earthworms after 21-d exposure ranged from below detection to 127 mg kg⁻¹ wet weight with the rank order of PFNA > PFHxS > PFHpA > PFBS; no mortality of earthworms was

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

	observed in all treatments including controls, except PFBS at 1,000 µg kg ⁻¹ and all PFASs at 100,000 µg kg ⁻¹ . The highest weight loss (29%) was observed for earthworms exposed to PFNA at 100,000 µg kg ⁻¹ , which was significantly different from all other treatments except PFHpA at 100,000 µg kg ⁻¹ (Karnjanapiboonwong et al 2018). ⁴⁸
<i>Aquatic Toxicity: LC₅₀, EC₅₀, ErC₅₀, NOAEC/NOEC</i>	<p>EC₅₀; Species: <i>Geitlerinema amphibium</i> (Blue-green Algae) Exponential Growth Phase BA-13; Conditions: saltwater, static, 20 deg C, pH 7.6-7.8; Concentration: 1.42 mM for 72 hr; Effect: decreased population growth /formulation/⁴⁹</p> <p>EC₅₀; Species: <i>Skeletonema marinoi</i> (Diatom) Exponential Growth Phase BA-98; Conditions: saltwater, static, 20 deg C, pH 7.6-7.8; Concentration: 2.4 mM for 72 hr; Effect: decreased population growth /formulation/⁵⁰</p> <p>EC₅₀; Species: <i>Chlorella vulgaris</i> (Green Algae) Exponential Growth Phase BA-02; Conditions: saltwater, static, 20 deg C, pH 7.6-7.8; Concentration: 5.21 mM for 72 hr; Effect: decreased population growth /formulation/⁵¹</p> <p>LC₅₀ = 108.6 mg/L/48H (other fish)⁵²</p>
<i>Mammalian Toxicity: LC₅₀, EC₅₀, ErC₅₀, NOAEC/NOEC</i>	
<i>Wildlife Toxicity: LC₅₀, EC₅₀, ErC₅₀, NOAEC/NOEC</i>	
<i>Breakdown/degradation /combustion products</i>	
<i>Anaerobic degradation</i>	
<i>Aerobic degradation</i>	
<i>Other observable ecological effects (e.g. BOD)</i>	
<i>Fate and Transport: Aquatic</i>	<p>"Of the PFCs measured, PFOS, PFHxS, PFBS, PFPrS, PFEtS, PFOSA, N-EtFOSAA, PFDoDA, PFUnDA, PFDA, PFNA, PFHpA, PFHxA, PFPeA, PFBA, and PFPrA were found at detectable concentrations in the tap water samples. The water samples from Shanghai (China) contained the greatest concentrations of total PFCs (arithmetic mean = 130 ng/L), whereas those from Toyama (Japan) contained only 0.62 ng/L (Mak et al 2009)."⁵³</p> <p>"In tap water, PFOS and PFOA levels ranged between 0.39 and 0.87 ng/L (0.78 and 1.74 pmol/L) and between 0.32 and 6.28 ng/L (0.77 and 15.2 pmol/L), respectively. PFHpA, PFHxS, and PFNA were also other detected PFCs. PFC levels were notably lower in bottled water, where PFOS could not be detected in any sample. Moreover, PFHpA, PFHxS,</p>

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

	PFOA, PFNA, PFOS, PFOSA, and PFDA could be detected in the river water samples. PFOS and PFOA concentrations were between <0.24 and 5.88 ng/L (<0.48 and 11.8 pmol/L) and between <0.22 and 24.9 ng/L (<0.53 and 60.1 pmol/L), respectively (Ericson 2008). ⁵⁴
<i>Fate and Transport: Terrestrial</i>	
<i>Fate and Transport: Atmospheric</i>	
<i>Transport Issues</i>	
<i>Factors affecting bioavailability</i>	
Global Environmental Impacts	
<i>Ozone Depletion Potential (ODP)</i>	
<i>Global Climate Change</i>	
<i>Greenhouse Gas Production</i>	
<i>Acid Rain Formation</i>	
Special Reports	
<i>EU</i>	<p>Short-chain Polyfluoroalkyl Substances (PFAS) – A literature review of information on human health effects and environmental fate and effect aspects of short-chain PFAS, Environmental project No. 1707, 2015 http://www2.mst.dk/Udgiv/publications/2015/05/978-87-93352-15-5.pdf</p> <p>Polyfluoroalkyl substances (PFASs) in textiles for children – Survey of chemical substances in consumer products No. 136, 2015 http://www2.mst.dk/Udgiv/publications/2015/04/978-87-93352-12-4.pdf</p> <p>Survey of PFOS, PFOA and other perfluoroalkyl and polyfluoroalkyl substances – Part of the LOUS-review, Environmental project No. 1475, 2013 http://www2.mst.dk/Udgiv/publications/2013/04/978-87-93026-03-2.pdf</p>

Notes on chemical research: Not found in NIOSH Pocket Guide; CCRIS; GENETOX; Haz-Map

¹ www.expub.com; Chemical Identity Page for Perfluoroheptanoic acid.

² www.expub.com; RTECS for Perfluoroheptanoic acid.

³ www.expub.com; Chemical Identity Page for Perfluoroheptanoic acid.

⁴ U.S. National Library of Medicine, ChemIDplus, a Toxnet Database, entry for "Perfluoroheptanoic acid.", accessed online at: <https://chem.sis.nlm.nih.gov/chemidplus/rn/startswith/375-85-9>

⁵ U.S. National Library of Medicine, ChemIDplus, a Toxnet Database, entry for "Perfluoroheptanoic acid.", accessed online at: <https://chem.sis.nlm.nih.gov/chemidplus/rn/startswith/375-85-9>

⁶ **HSDB 2018a**: Hazardous Substances Data Bank. Bethesda (MD): National Library of Medicine (US); Last Revision Date October 25, 2016; cited 2018 Jun. Perfluoroheptanoic acid, CASRN: 375-85-9; Hazardous Substances Databank Number: 8293. Available from: <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

Siegemund G et al; Fluorine Compounds, Organic. Ullmann's Encyclopedia of Industrial Chemistry 7th ed. (1999-2015). NY, NY: John Wiley & Sons. Online Posting Date: June 15, 2000] **PEER REVIEWED**

⁷ HSDB 2018a: [Sigma-Aldrich; Product Search Perfluoroheptanoic Acid (375-85-9). Available from, as of Jan 6, 2016: <http://www.sigmaaldrich.com/united-states.html>] **PEER REVIEWED**

⁸ HSDB 2018a: [Royal Soc Chem; ChemSpider. Perfluoroheptanoic acid. (375-85-9). Available from, as of Dec 15, 2015: <http://www.chemspider.com/Search.aspx>] **PEER REVIEWED**

⁹ HSDB 2018a: [Sigma-Aldrich; Product Search Perfluoroheptanoic Acid (375-85-9). Available from, as of Jan 6, 2016: <http://www.sigmaaldrich.com/united-states.html>] **PEER REVIEWED**

¹⁰ HSDB 2018a: [Siegemund G et al; Fluorine Compounds, Organic. Ullmann's Encyclopedia of Industrial Chemistry. 7th ed. (1999-2015). New York, NY: John Wiley & Sons. Online Posting Date: 15 Jun 2000] **PEER REVIEWED**

¹¹ HSDB 2018a: [US EPA; Estimation Program Interface (EPI) Suite. Ver. 4.11. Nov, 2012. Available from, as of Dec 15, 2015: <http://www2.epa.gov/tsca-screening-tools>] **PEER REVIEWED**

¹² HSDB 2018a: [Siegemund G et al; Fluorine Compounds, Organic. Ullmann's Encyclopedia of Industrial Chemistry. 7th ed. (1999-2015). New York, NY: John Wiley & Sons. Online Posting Date: 15 Jun 2000] **PEER REVIEWED**

¹³ HSDB 2018a: [Steele WV et al; *J Chem Eng Data* **47**: 648-66 (2002)] **PEER REVIEWED**

¹⁴ HSDB 2018a: [Sigma-Aldrich; Safety Data Sheet for Perfluoroheptanoic acid. Product Number: 342041, Version 4.14 (Revision Date 06/27/2014). Available from, as of December 15, 2015: <http://www.sigmaaldrich.com/safety-center.html>] **PEER REVIEWED**

¹⁵ HSDB 2018a: [Sigma-Aldrich; Safety Data Sheet for Perfluoroheptanoic acid. Product Number: 342041, Version 4.14 (Revision Date 06/27/2014). Available from, as of December 15, 2015: <http://www.sigmaaldrich.com/safety-center.html>] **PEER REVIEWED**

¹⁶ HSDB 2018a: [Royal Soc Chem; ChemSpider. Perfluoroheptanoic acid. (375-85-9). Available from, as of Dec 15, 2015: <http://www.chemspider.com/Search.aspx>] **PEER REVIEWED**

¹⁷ HSDB 2018a: [Sigma-Aldrich; Safety Data Sheet for Perfluoroheptanoic acid. Product Number: 342041, Version 4.14 (Revision Date 06/27/2014). Available from, as of December 15, 2015: <http://www.sigmaaldrich.com/safety-center.html>] **PEER REVIEWED**

¹⁸ **NICNAS 2018**: Australian Government, Department of Health, National Industrial Chemicals Notification and Assessment Scheme (NICNAS). HUMAN HEALTH TIER II ASSESSMENT FOR Perfluoroheptanoic acid (PFHpA) and its direct precursors. Accessed online at: https://www.nicnas.gov.au/chemical-information/imap-assessments/imap-group-assessment-report?assessment_id=1689.

¹⁹ **Wang 2017**: Wang B, et al. Perfluoroalkyl substances and endometriosis-related infertility in Chinese women. *Environ Int.* 2017 May; **102**:207-212.

²⁰ **Mulkiewicz 2007**: Mulkiewicz E, et al. Evaluation of the acute toxicity of perfluorinated carboxylic acids using eukaryotic cell lines, bacteria and enzymatic assays. *Environmental Toxicology and Pharmacology* **23** (2007) 279-285.

²¹ **Kleszczynski 2007**: Kleszczynski K, et al. Analysis of structure-cytotoxicity in vitro relationship (SAR) for perfluorinated carboxylic acids. *Toxicology in Vitro* **21** (2007) 1206-1211.

²² **Nobels 2010**: Nobels I, et al. Application of a multiple endpoint bacterial reporter assay to evaluate toxicological relevant endpoints of perfluorinated compounds with different functional groups and varying chain length. *Toxicol In Vitro.* 2010 Sept; **24**(6): 1768-74.

²³ **NICNAS 2018**, under section on Carcinogenicity; **Wolf 2012**: Wolf CJ, et al. Activation of mouse and human peroxisome-proliferator-activated receptor-alpha (PPAR α) by perfluoroalkyl acids (PFAAs): Further investigation of C4-C12 compounds. *Reproductive Toxicology* **33**(4) July 2012, 546-551.

²⁴ **NICNAS 2018**, under section on Repeated Dose Toxicity – Oral.

²⁵ **NICNAS 2018**, under section on Corrosion/Irritation – Skin Irritation.

²⁶ HSDB 2018a: [Royal Soc Chem; ChemSpider. Perfluoroheptanoic acid. (375-85-9). Available from, as of Dec 15, 2015: <http://www.chemspider.com/Search.aspx>] **PEER REVIEWED**

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

- ²⁷ **Ruan 2014:** Ruan T, et al. 6:2 Fluorotelomer iodide in vitro metabolism by rat liver microsomes: comparison with [1,2-(14)C] 6:2 fluorotelomer alcohol. *Chemosphere*. 2014 Oct; **112**:34-41.
- ²⁸ **Russell 2015:** Russell MH, et al. Inhalation and oral toxicokinetics of 6:2 FTOH and its metabolites in mammals. *Chemosphere*. 2015 Feb; **120**:328-35.
- ²⁹ **Fujii 2015:** Fujii, Y, et al. Toxicokinetics of perfluoroalkyl carboxylic acids with different carbon chain lengths in mice and humans. *Journal of Occupational Health* 2015; **57**: 1-12.
- ³⁰ NICNAS 2018, under section on Toxicokinetics.
- ³¹ NICNAS 2018, under section on Toxicokinetics.
- ³² **CT DPH 2017:** Connecticut Department of Public Health – Environmental & Occupational Health Assessment Program. Perfluoroalkyl Substances (PFAS) in Drinking Water: Health Concerns Fact Sheet. October 2017. Accessed online at: http://portal.ct.gov/-/media/Departments-and-Agencies/DPH/dph/environmental_health/eoha/Toxicology_Risk_Assessment/2018-uploads/Perfluoroalkyl-Substances-PFASs-in-DWHealth-Concerns.pdf?la=en.
- ³³ HSDB 2018a: [40 CFR 141.40 (USEPA); U.S. National Archives and Records Administration's Electronic Code of Federal Regulations. Available from, as of December 14, 2015: <http://www.ecfr.gov>] **PEER REVIEWED**
- ³⁴ **ECHA 2018a:** European Chemicals Agency record for Perfluoroheptanoic acid, CAS # 375-85-9, Accessed online at: <https://echa.europa.eu/information-on-chemicals/cl-inventory-database/-/discli/details/37168>.
- ³⁵ **Knobeloch 2012:** Knobeloch L, et al. Perfluoroalkyl chemicals in vacuum cleaner dust from 39 Wisconsin homes. *Chemosphere*. 2012 Aug; **88**(7): 779-83.
- ³⁶ **Hu 2016:** Hu XC, et al. Detection of Poly- and Perfluoroalkyl Substances (PFASs) in U.S. Drinking Water Linked to Industrial Sites, Military Fire Training Areas, and Wastewater Treatment Plants. *Environmental Science and Technology Letters* 2016, **3**, 344-350.
- ³⁷ **Ng and Hungerbuhler 2014:** Ng, CA and Hungerbuhler K. Bioaccumulation of Perfluorinated Alkyl Acids: Observations and Models. *Environmental Science & Technology* 2014, **48**, 4637-4648.
- ³⁸ **D'Hollander 2010:** D'Hollander W, et al. Perfluorinated substances in human food and other sources of human exposure. *Rev Environ Contam Toxicol*. 2010; **208**: 179-215.
- ³⁹ **Wang 2016c:** Wang B, et al. Perfluoroalkyl and polyfluoroalkyl substances in cord blood of newborns in Shanghai, China: Implications for risk assessment. *Environment International* **97** (2016) 7-14.
- ⁴⁰ **ATSDR 2015:** ATSDR Draft Toxicological Profile for Perfluoroalkyls. Page 392. August 2015. Accessed online at: <https://www.atsdr.cdc.gov/toxprofiles/tp200.pdf>. Pages 411-413.
- ⁴¹ ATSDR 2015. Page 392.
- ⁴² **ECHA 2015a:** European Chemicals Agency, (August 31, 2015). DECISION ON SUBSTANCE EVALUATION PURSUANT TO ARTICLE 46(1) OF REGULATION (EC) NO 1907/2006 – For reaction mass of mixed (3,3,4,4,5,5,6,6,7,7,8,8,8-tridecafluorooctyl) phosphates, ammonium salt, CAS No not available (EC No 700-161-3). Accessed online: <https://echa.europa.eu/documents/10162/58786c6a-d440-46d7-9438-022fdc37191c>. Page 4.
- ⁴³ **Zhao 2013:** Zhao S, et al. Bioaccumulation of perfluoroalkyl carboxylates (PFCAs) and perfluoroalkane sulfonates (PFASs) by earthworms (*Eisenia fetida*) in soil. *Environmental Pollution* **179** (2013) 45-52.
- ⁴⁴ **Zhang 2013:** Zhang T, et al. Distribution of Poly- and Perfluoroalkyl Substances in Matched Samples from Pregnant Women and Carbon Chain Length Related Maternal Transfer. *Environmental Science & Technology*, 2013, **47**(14), 7974-7981.
- ⁴⁵ **Dalahmeh 2018:** Dalahmeh S, et al. Per- and polyfluoroalkyl substances (PFASs) in water, soil and plants in wetlands and agricultural areas in Kampala, Uganda. *Science of the Total Environment* **631-632** (2018) 660-667.
- ⁴⁶ **Ishibashi 2011:** Ishibashi H, et al. Transactivation potencies of the Baikal seal (*Pusa sibirica*) peroxisome proliferator—activated receptor α by perfluoroalkyl carboxylates and sulfonates: estimation of PFOA induction equivalency factors. *Environmental Science & Technology* 2011 Apr 1; **45**(7): 3123-30.

Updated EHS Summary of Perfluoroheptanoic acid for the MA TURA Science Advisory Board Meeting – June 21, 2018

- ⁴⁷ **Kim 2015:** Kim M, et al. Perfluoroheptanoic acid affects amphibian embryogenesis by inducing the phosphorylation of ERK and JNK. *International Journal of Molecular Medicine*, December 2015, **36 (6)**, 1693-1700.
- ⁴⁸ **Karnjanapiboonwong 2018:** Karnjanapiboonwong A, et al. Perfluoroalkylsulfonic and carboxylic acids in earthworms (*Eisenia fetida*): Accumulation and effects results from spiked soils at PFAS concentrations bracketing environmental relevance. *Chemosphere*. 2018 May; **199**: 168-173.
- ⁴⁹ **HSDB 2018a:** [Latala A, et al; *Environ Toxicol Pharmacol* **28(2)**: 167-71 (2009) as cited in the ECOTOX database. Available from, as of February 22, 2016: http://cfpub.epa.gov/ecotox/quick_query.htm] **PEER REVIEWED**
- ⁵⁰ HSDB 2018a.
- ⁵¹ HSDB 2018a.
- ⁵² www.expub.com; RTECS for Perfluoroheptanoic acid.
- ⁵³ **Mak 2009:** Mak YL, et al. Perfluorinated compounds in tap water from China and several other countries. *Environmental Science & Technology* 2009 Jul 1; **43(13)**: 4824-9.
- ⁵⁴ **Ericson 2008:** Ericson I, et al. Levels of perfluorochemicals in water samples from Catalonia, Spain: is drinking water a significant contribution to human exposure? *Environmental Science and Pollution Research Int.* 2008 Oct; **15(7)**: 614-9.